



BEP – 5 Days

This leaflet is offered as a guide to you and your family. The possible benefits of treatment vary; for some people chemotherapy may reduce the risk of the cancer coming back, for others it may control the cancer and its symptoms. Chemotherapy is the most commonly prescribed anti-cancer treatment but other types of treatment are also used. Your doctor will explain to you whether you will receive chemotherapy or another type of treatment, or a combination of both. Your doctor or nurse will be happy to answer any questions you have about your treatment. You will find it useful to refer to the booklet *Chemotherapy, a guide* which gives general information on chemotherapy and side effects.

Your treatment

Your doctor has prescribed for you a treatment known as 5 Day BEP.

Your treatment consists of the following:

| | |
|--------------------------|---|
| Days 1 to 5 | Etoposide and Cisplatin via a drip over 1 hours |
| Days 2, 12 and 19 | Bleomycin via a drip over 30 minutes |

In addition to the chemotherapy we will also give you intravenous hydration (fluids).

This is repeated every 3 weeks for 3 or 4 cycles.

You will have a routine blood test before the start of each cycle of treatment.

This treatment can have serious or possibly life-threatening side effects. It is very important that you report side effects straight away. Don't delay, if you feel unwell, please ring The Christie Hotline on 0161 446 3658. The lines are open 24 hours a day.

Increased risk of serious infection:

You are vulnerable to infection while you are having chemotherapy. Minor infections can become life-threatening in a matter of hours if left untreated. Symptoms of infection include fever, shivering, sweats, sore throat, diarrhoea, discomfort when you pass urine, cough or breathlessness. We recommend that you use a digital thermometer so you can check your temperature. You can buy one from your local chemist.

If your temperature is 37.5°C or above or below 36 °C or you have symptoms of an infection, contact The Christie Hotline straight away.

Possible side effects

Chemotherapy can cause many different side effects. Some are more likely to occur than others. Everyone is different and not everyone gets all the side effects. Most side effects are usually temporary, but in some rare cases they can be life-threatening. It is important to tell your hospital doctor or nurse about any side effects so they can be monitored and, where possible, treated.

Common side effects (more than 1 in 10)

- **Anaemia (low number of red blood cells)**

While having this treatment you may become anaemic. This may make you feel tired and breathless. Let your doctor or nurse know if these symptoms are a problem. You may need a blood transfusion.

- **Bruising or bleeding**

This treatment can reduce the production of platelets which help the blood clot. Let your doctor know if you have any unexplained bruising or bleeding, such as nosebleeds, bloodspots or rashes on the skin, and bleeding gums. You may need a platelet transfusion.

- **Nausea and vomiting (sickness)**

The severity of this varies from person to person. Anti-sickness medication may be given along with your chemotherapy to prevent this. You may also be given anti-sickness tablets to take at home. If you continue to feel or be sick, contact your GP or this hospital, because your anti-sickness medication may need to be changed or increased.

- **Lethargy**

Some chemotherapy may make you feel tired and lacking in energy. It can be frustrating when you feel unable to cope with routine tasks. If you do feel tired, take rest and get help with household chores. If necessary, take time off work. Gentle exercise such as walking can be beneficial.

- **Tinnitus & high frequency hearing loss**

You may develop tinnitus (ringing in the ears), this sensation should subside when your treatment finishes. Please tell your doctor if this occurs. High frequency hearing loss can also occur with this chemotherapy, this may be permanent.

- **Hair loss**

Hair loss is usually total. The hair falls out gradually 10 to 14 days following your first course of treatment. The time scale varies from person to person. Please remember that this is a temporary side effect and your hair will grow back when your treatment is completed. If you would like an appointment with the wig service, this can be arranged for you. Ask the staff for a copy of the *Wig Fitting Service* leaflet.

- **Sore mouth**

Your mouth may become sore or dry, or you may notice small mouth ulcers during this treatment. Drinking plenty of fluids and cleaning your teeth regularly and gently with a soft toothbrush can help to reduce the risk of this happening. We can prescribe a mouthwash for you to use during treatment. You can dilute this with water if your mouth is sore. Ask your doctor or nurse for further advice. There is also general mouth care information in the chemotherapy booklet. If you continue to have a sore mouth, please contact The Christie Hotline.

- **Diarrhoea**

If this becomes a problem while you are having treatment, anti-diarrhoea tablets can be bought from a pharmacy or prescribed by your GP for a temporary period until this is resolved. If the problem persists contact this hospital. **If you develop severe diarrhoea it is important to contact this hospital straightaway as this may be a sign of a serious infection. Don't delay!**

- **Strange taste**

Occasionally during treatment you may experience a strange taste, sometimes described as metallic or bitter. A strongly flavoured sweet or mint will help to disguise this.

- **Skin changes**

If the skin on your **hands and feet becomes sore and fingertips dry and cracked**, contact the hospital straightaway. Tablets and cream can be prescribed to help you. Also, try and keep your hands and feet cool and if possible uncovered, or wear loose-fitting cotton socks. You may also develop **ridging of the nails**, this will go when treatment finishes. Some chemotherapy can make **your skin more sensitive to the sun** than usual. Sit in the shade, avoid too much sun and use a high factor sunblock cream. Asian and African-Caribbean people may develop noticeable light patches on their skin.

- **Tingling & numbness in the fingers or toes**

This is only usually mild and temporary but sometimes lasts for some time or may become permanent. Please report these symptoms to your doctor on your next hospital visit

- **Flu-like symptoms**

Bleomycin may cause flu-like symptoms such as fever, aches and pains and shivering about 3 to 5 hours after it is given. These symptoms are temporary and should go within 12 to 24 hours. Paracetamol will help. If your symptoms are particularly severe, tell your doctor on your next visit.

- **Kidney function**

Some chemotherapy such as cisplatin can affect your kidneys. It is important to monitor how your kidneys are working while you are having treatment.

Uncommon side effects (less than 1 in 10)



- **Lung function**

Bleomycin may cause some changes to your lung tissue. You may not be aware of these changes but because of this potential side effect, we will test your lung function both before and after your treatment. **Please report any cough or breathlessness to your doctor without delay.**

- **Herbal medicine**

Some herbal medicine including St John's Wort can affect the chemotherapy. You should let your doctor or nurse know if you are taking any herbal medication, complementary or alternative medicines, including vitamins, minerals and medicines purchased over-the-counter.

- **Extravasation**

Extravasation is when chemotherapy leaks outside the vein. If you develop redness, soreness or pain at the injection site **at any time** please let us know straightaway.

Serious and potentially life threatening side effects

In a small proportion of patients chemotherapy can result in very severe side effects which may rarely result in death. The team caring for you will discuss the risk of these side effects with you.

Sex, contraception & fertility

Protecting your partner and contraception: We recommend that you or your partner use a condom during sexual intercourse while you are having the course of chemotherapy. Chemotherapy is dangerous to unborn babies and this will also protect you and your partner from any chemotherapy drugs that may be present in semen and in the vagina. If you suspect that you may be pregnant please tell your doctor immediately.

Fertility: This chemotherapy may affect your ability to have children. Your doctor or nurse should have discussed this with you. If not, please ask them before you start treatment.

Late side effects

Some side effects may become evident only after a number of years. In reaching any decision with you about treatment, the potential benefit you receive from treatment will be weighed against the risks of serious long term side effects to the heart, lungs, kidneys and bone marrow. With some drugs there is also a small but definite risk of developing another cancer. If any of these problems specifically applies to you, the doctor will discuss these with you and note this on your consent form.

