



# Dacarbazine

This leaflet is offered as a guide to you and your family. The possible benefits of treatment vary: for some people chemotherapy may reduce the risk of the cancer coming back, for others chemotherapy may control the cancer and its symptoms. Chemotherapy is the most commonly prescribed anti-cancer treatment but other types of treatment are also used. Your doctor will explain to you whether you will receive chemotherapy or another type of treatment, or a combination of both. Your doctor or nurse will be happy to answer any questions. You will find it useful to refer to the booklet *Chemotherapy: a guide* which gives general information on chemotherapy and side effects.

## Your treatment

Your doctor or nurse clinician has prescribed for you a regimen of treatment known as dacarbazine or DTIC.

**Day 1** Dacarbazine is given as a intravenous infusion (drip) which lasts for 30 minutes, but can be given over 60 minutes if vein pain is a problem.

**You will have treatment every 3 weeks for up to 6 cycles.**

You will have a routine blood test before the start of each cycle of treatment.

This treatment can have serious or possibly life-threatening side effects. It is very important that you report side effects straight away. Don't delay, if you feel unwell, please ring The Christie Hotline on 0161 446 3658. The lines are open 24 hours a day.

## Possible side effects

Chemotherapy can cause many different side effects. Some are more likely to occur than others. Everyone is different and not everyone gets all the side effects. Most side effects are usually temporary, but in some rare cases they can be life-threatening. It is important to tell your hospital doctor or nurse about any side effects so they can be monitored and, where possible, treated.

### Common side effects (more than 1 in 10)

#### Increased risk of serious infection:

You are vulnerable to infection while you are having chemotherapy. Minor infections can become life-threatening in a matter of hours if left untreated. Symptoms of infection include fever, shivering, sweats, sore throat, diarrhoea, discomfort when you pass urine, cough or breathlessness. We recommend that you use a digital thermometer so you can check your temperature. You can buy one from your local chemist.

**If you feel unwell, you have symptoms of an infection or your temperature is 37.5°C or above or below 36°C contact The Christie Hotline straight away.**

- **Vein pain**

This chemotherapy can cause pain along the vein during and after treatment. This should only be temporary but contact your hospital doctor or nurse if this becomes severe.

- **Loss of appetite**

Your appetite may be reduced. A dietitian or specialist nurse at The Christie can give you advice and tips on boosting appetite, coping with eating difficulties and maintaining weight. Ask staff for a copy of 'Eating – help yourself'.

- **Nausea and vomiting (sickness)**

The severity of this varies from person to person. Anti-sickness medication may be given along with your chemotherapy to prevent this. You may also be given anti-sickness tablets to take at home. If you continue to feel or be sick, contact your GP or The Christie, because your anti-sickness medication may need to be changed or increased.

### **Uncommon side effects (less than 1 in 10)**

- **Flu-like symptoms**

Some chemotherapy may cause flu-like symptoms such as fever, aches and pains and shivering about 3 to 5 hours after it is given. These symptoms are temporary and should go within 12 to 24 hours. Paracetamol will help. If your symptoms are particularly severe, tell your doctor on your next visit.

- **Hair loss**

Some hair loss may occur during treatment. It is advisable to avoid perms, colours, use of hot brushes and vigorous, frequent washing that could increase hair loss. Please remember that this is a temporary side effect and your hair will grow back when your treatment is completed. If you would like an appointment with the wig service, this can be arranged for you. Ask the staff for a copy of the 'Wig Fitting Service at The Christie'.

- **Sensitivity to the sun**

Some chemotherapy can make your skin more sensitive to the sun than usual. Sit in the shade, avoid too much sun and use a sunblock cream.

- **Hyperpigmentation**

Your skin may appear darker in colour or lightly tanned, especially around the joints. This is known as hyperpigmentation. Asian and African-Caribbean people may develop noticeable light patches on their skin. The skin will return to normal when treatment is finished.

- **Lethargy**

Some chemotherapy may make you feel tired and lacking in energy. It can be frustrating when you feel unable to cope with routine tasks. If you do feel tired, take rest and get help with household chores. If necessary, take time off work. Gentle exercise such as walking can be beneficial.

## Rare side effects (less than 1 in 100)

- **Immediate allergic reactions:**

Ask the staff for help if you notice any of the following:

- Some people have hot flushes when the drug is being given.
- Fevers and chills: back pain, shortness of breath, headaches and swelling of the face may occur *during* the time the drug is being given. If this happens please tell the staff straightaway. Your doctor may prescribe a drug that can help to reduce these side effects and, if necessary, we can give you this before your next treatment.

- **Anaemia (low number of red blood cells)**

While having this treatment you may become anaemic. This may make you feel tired and breathless. Let your doctor or nurse know if these symptoms are a problem. You may need a blood transfusion.

- **Bruising or bleeding**

This treatment can reduce the production of platelets which help the blood clot. Let your doctor know if you have any unexplained bruising or bleeding, such as nosebleeds, bloodspots or rashes on the skin, and bleeding gums. You may need a platelet transfusion.

- **Diarrhoea**

If this becomes a problem while you are having treatment, anti-diarrhoea medication can be prescribed by your GP for a temporary period until this is resolved. If the problem persists contact this hospital. **If you develop severe diarrhoea it is important to contact this hospital straight away as this may be a sign of a serious infection. Don't delay!**

- **Extravasation** is when chemotherapy leaks outside the vein. If you develop redness, soreness or pain at the injection site **at any time** please let us know straightaway.

- **Kidney function**

Some chemotherapy can affect your kidneys. We will monitor how your kidneys are working while you are having treatment.

## Serious and potentially life threatening side effects

In a small proportion of patients chemotherapy can result in very severe side effects which may rarely result in death. The team caring for you will discuss the risk of these side effects with you.

## Sex, contraception & fertility

**Protecting your partner and contraception:** We recommend that you or your partner use a condom during sexual intercourse while you are having the course of chemotherapy. Chemotherapy is dangerous to unborn babies and this will also protect you and your partner from any chemotherapy drugs that may be present in semen and in the vagina. If you suspect that you may be pregnant please tell your doctor immediately.

**Fertility:** This chemotherapy may affect your ability to have children. Your doctor or nurse should have discussed this with you. If not, please ask them before you start treatment.

## Late side effects

Some side effects may become evident only after a number of years. In reaching any decision with you about treatment, the potential benefit you receive from treatment will be weighed against the risks of serious long term side effects to the heart, lungs, kidneys and bone marrow. With some drugs there is also a small but definite risk of developing another cancer. If any of these problems specifically applies to you, the doctor will discuss these with you and note this on your consent form.

## Contacts

If you have any general questions or concerns about your treatment, please ring the area where you are having treatment:

Contacts:

- Administration enquiries 0161 918 7606/7610
- Chemotherapy nurse: 0161 918 7171
- Clinical trials unit 0161 918 7663

For urgent advice ring The Christie Hotline on 0161 446 3658 (24 hours)

Your consultant is: .....

Your hospital number is: .....

Your key worker is: .....



**The Christie Patient Information Service July 2014**  
**CHR/CT/257/12.01.04 Version 5 Review date: July 2017**

Details of the sources used are available, please contact [Patient.Information@christie.nhs.uk](mailto:Patient.Information@christie.nhs.uk)

The Christie Hotline 0161 446 3658