



# Mitomycin C and Fluorouracil

This leaflet is offered as a guide to you and your family. The possible benefits of treatment vary; for some people chemotherapy may reduce the risk of the cancer coming back, for others it may control the cancer and its symptoms. Chemotherapy is the most commonly prescribed anti-cancer treatment but other types of treatment are also used. Your doctor will explain to you whether you will receive chemotherapy or another type of treatment, or a combination of both. Your doctor or nurse will be happy to answer any questions you have about your treatment. You will find it useful to refer to the booklet *Chemotherapy, a guide* which gives general information on chemotherapy and side effects.

## Your treatment

Your doctor has prescribed for you a treatment which includes the chemotherapy Mitomycin C and Fluorouracil, also called 5FU. Your treatment will be given:

**Day 1:** Mitomycin C by bolus (short injection) followed by  
Fluorouracil given via a drip over ..... or  
Via a pump over ..... days.

The treatment is repeated every ..... weeks for ..... cycles.

You will have a routine blood test before the start of each cycle of treatment.

**Extravasation** is when chemotherapy leaks outside the vein. If you develop redness, soreness or pain at the injection site **at any time** please let us know straight away.

This treatment can have serious or possibly life-threatening side effects. It is very important that you report side effects straight away. Don't delay, if you feel unwell, please ring The Christie Hotline on 0161 446 3658. The lines are open 24 hours a day.

## Increased risk of serious infection

You are vulnerable to infection while you are having chemotherapy. Minor infections can become life-threatening in a matter of hours if left untreated. Symptoms of infection include fever, shivering, sweats, sore throat, diarrhoea, discomfort when you pass urine, cough or breathlessness. We recommend that you use a digital thermometer so you can check your temperature. You can buy one from your local chemist.

**If you feel unwell, you have symptoms of an infection or your temperature is 37.5°C or above or below 36°C contact The Christie Hotline straight away.**

## Possible side effects

Chemotherapy can cause many different side effects. Some are more likely to occur than others. Everyone is different and not everyone gets all the side effects. Most side effects are usually temporary, but in some rare cases they can be life-threatening. It is important to tell your hospital doctor or nurse about any side effects so they can be monitored and, where possible, treated.



- **Chest pain or stroke**

A small number of patients receiving 5-Fluorouracil (5FU) or capecitabine can experience chest pain (angina) or rarely have a heart attack. Extremely rarely this may lead to death. Other complications such as stroke or mini-stroke can happen but are exceptionally rare. If you develop any of these symptoms you should contact your hospital doctor for advice. In an emergency you should go immediately to your nearest accident and emergency department.

- **Anaemia (low number of red blood cells)**

While having this treatment you may become anaemic. This may make you feel tired and breathless. Let your doctor or nurse know if these symptoms are a problem. You may need a blood transfusion.



- **Bruising or bleeding**

This treatment can reduce the production of platelets which help the blood clot. Let your doctor know if you have any unexplained bruising or bleeding, such as nosebleeds, bloodspots or rashes on the skin, and bleeding gums. You may need a platelet transfusion.

- **Nausea and vomiting (sickness)**

The severity of this varies from person to person. Anti-sickness medication may be given along with your chemotherapy to prevent this. You may also be given anti-sickness tablets to take at home. If you continue to feel or be sick, contact your GP or this hospital, because your anti-sickness medication may need to be changed or increased.



- **Diarrhoea**

If this becomes a problem while you are having treatment, anti-diarrhoea tablets can be bought from a pharmacy or prescribed by your GP for a temporary period until this is resolved. If the problem persists contact this hospital. Ask the staff for a copy of *Eating: Help yourself* which has some useful ideas about diet when you are having treatment.

- **Lethargy**

Some chemotherapy may make you feel tired and lacking in energy. It can be frustrating when you feel unable to cope with routine tasks. If you do feel tired, take rest and get help with household chores. If necessary, take time off work. Gentle exercise such as walking can be beneficial.

- **Watery eyes**

Your eyes may also water. Often, this will improve in time and needs no specific treatment but if you have ongoing symptoms please discuss this with your doctor or nurse.

- **Sore mouth**

Your mouth may become sore or dry, or you may notice small mouth ulcers during this treatment. Drinking plenty of fluids and cleaning your teeth regularly and gently with a soft toothbrush can help to reduce the risk of this happening. We can prescribe a mouthwash for you to use during treatment. You can dilute this with water if your mouth is sore. Ask your doctor or nurse for further advice. There is also general mouth care information in the chemotherapy booklet. If you continue to have a sore mouth, please contact The Christie Hotline.

- **Skin and nail changes**

**PPE (palmar-plantar erythema):** The skin on your hands and feet may become very dry, red and sore with some cracking. Tell your doctor. Cream and tablets can be prescribed to help. Your chemotherapy dose may need to change. Try to keep your hands and feet cool and if possible, uncovered.

**Hyperpigmentation:** Your skin may appear darker in colour or lightly tanned, especially around the joints. This is known as hyperpigmentation. Asian and African-Caribbean people may develop noticeable light patches on their skin. The skin will return to normal when treatment is finished.

**Increased sensitivity to the sun:** Your skin will tan or burn in the sun more easily. Sit in the shade, avoid too much sun and use a high factor sunblock cream and wear a hat.

- **Darkening of the veins**

Fluorouracil can irritate the vein wall and can cause discolouration. So you may get brown marks along the course of the vein where the drip was inserted, and your arm can also become itchy. This will usually fade over time. Please tell your hospital doctor at your next appointment if you have any problems.

- **Hair thinning**

Some hair loss may occur during treatment although this is unlikely. It is advisable to avoid perms, colours, use of hot brushes and vigorous, frequent washing that could increase hair loss. Please remember that this is a temporary side effect and your hair will grow back when your treatment is completed. If you would like an appointment with the wig service, this can be arranged for you. Ask the staff for a copy of *'The Wig Fitting Service.'*

## **Rare side effects (less than 1 in 100)**

### **Serious and potentially life-threatening side effects**

In a small proportion of patients chemotherapy can result in very severe side effects which may rarely result in death. The team caring for you will discuss the risk of these side effects with you.

## Sex, contraception & fertility

**Protecting your partner & contraception:** We recommend that you use a condom during sexual intercourse. Chemotherapy is dangerous to unborn babies and this will also protect you and your partner from any chemotherapy drugs that may be present in semen and in the vagina. If you suspect that you may be pregnant please tell your doctor immediately.

**Fertility:** This chemotherapy may affect your ability to have children. Your doctor or nurse should have discussed this with you. If not, please ask them before you start treatment.

## Late side effects

Some side effects may become evident only after a number of years. In reaching any decision with you about treatment, the potential benefit you receive from treatment will be weighed against the risks of serious long term side effects to the heart, lungs, kidneys and bone marrow. With some drugs there is also a small but definite risk of an increased chance of developing another cancer. If any of these problems specifically applies to you, the doctor will discuss these with you and note this on your consent form.

## Contacts

If you have any general questions or concerns about your treatment, please ring the area where you are having treatment:

- Administration enquiries 0161 918 7606/7610
- Chemotherapy nurse: 0161 918 7171
- Clinical trials unit 0161 918 7663

For advice ring The Christie Hotline on 0161 446 3658 (24 hours)

Your consultant is: .....

Your hospital number is: .....

Your key worker is: .....



**The Christie Patient Information Service July 2014**  
**CHR/CT/287/12.1.04 Version 5 Review date: July 2017**

Details of the sources used are available, please contact [Patient.Information@christie.nhs.uk](mailto:Patient.Information@christie.nhs.uk)

The Christie Hotline 0161 446 3658