



Irinotecan and Temozolomide

This leaflet is offered as a guide to you and your family. The possible benefits of treatment vary; for some people chemotherapy may reduce the risk of the cancer coming back, for others it may control the cancer and its symptoms. Chemotherapy is the most commonly prescribed anti-cancer treatment but other types of treatment are also used. Your doctor will explain to you whether you will receive chemotherapy or another type of treatment, or a combination of both. Your doctor or nurse will be happy to answer any questions you have about your treatment. You will find it useful to refer to the booklet *Chemotherapy, a guide* which gives general information on chemotherapy and side effects.

Your treatment

Your doctor or nurse clinician has prescribed for you a treatment which includes the chemotherapy irinotecan and temozolomide.

Day 1 to 5	Temozolomide capsules taken by mouth Irinotecan intravenous (IV) infusion over 1 hour
Day 6 and 7	No treatment
Day 8 to 12	Irinotecan IV infusion over 1 hour
Day 22	Restart as per day 1

Temozolomide should be taken on an empty stomach. We recommend that you do not eat for 1 to 2 hours before taking the temozolomide and for 1 to 2 hours afterwards.

Temozolomide capsules come in a variety of strengths (5mg, 20mg, 100mg, 250mg) so you may need to take several capsules to make up your daily dose.

You will have a routine blood test before the start of each cycle of treatment.

This treatment can have serious or possibly life-threatening side effects. It is very important that you report side effects straight away. Don't delay, if you feel unwell, please ring The Christie Hotline on 0161 446 3658. The lines are open 24 hours a day.

Increased risk of serious infection

You are vulnerable to infection while you are having chemotherapy. Minor infections can become life-threatening in a matter of hours if left untreated. Symptoms of infection include fever, shivering, sweats, sore throat, diarrhoea, discomfort when you pass urine, cough or breathlessness. We recommend that you use a digital thermometer so you can check your temperature. You can buy one from your local chemist.

If you feel unwell, you have symptoms of an infection or your temperature is 37.5°C or above or below 36 °C contact The Christie Hotline straight away.

Possible side effects

Chemotherapy can cause many different side effects. Some are more likely to occur than others. Everyone is different and not everyone gets all the side effects. Most side effects are usually temporary, but in some rare cases they can be life-threatening. It is important to tell your hospital doctor or nurse about any side effects so they can be monitored and, where possible, treated.

Common side effects (more than 1 in 10)

- **Nausea and vomiting (sickness)**

The severity of this varies from person to person. You will be given anti-sickness tablets to take home, and it is helpful to take an anti-sickness tablet around 30 minutes before your temozolomide tablets. If you continue to feel or be sick, contact your GP or The Christie because your anti-sickness medication may need to be changed or increased.

- **Hair thinning**

Some hair loss may occur during treatment although this is unlikely. It is advisable to avoid perms, colours, use of hot brushes and vigorous, frequent washing that could increase hair loss. Please remember that this is a temporary side effect and your hair will grow back when your treatment is completed. If you would like an appointment with the wig service, this can be arranged for you. Ask the staff for a copy of 'The Wig Fitting Service.'

Uncommon side effects (less than 1 in 10)

- **Lethargy**

Some chemotherapy may make you feel tired and lacking in energy. It can be frustrating when you feel unable to cope with routine tasks. If you do feel tired, take rest when necessary. Gentle exercise such as walking can be beneficial.

- **Anaemia (low number of red blood cells)**

While having this treatment you may become anaemic. This may make you feel tired and breathless. Let your doctor or nurse know if these symptoms are a problem. You may need a blood transfusion.

- **Bruising or bleeding**

This treatment can reduce the production of platelets which help the blood clot. Let your doctor know if you have any unexplained bruising or bleeding, such as nosebleeds, bloodspots or rashes on the skin, and bleeding gums. You may need a platelet transfusion.

- **Chemotherapy reaction**

Sometimes during the infusion or within 24 hours of the infusion of irinotecan you may get some of these symptoms: abdominal cramps, diarrhoea, sweats, dizziness, excess saliva, watery eyes, tiredness and occasionally problems with vision. These side effects can easily be managed with an injection underneath the skin. Tell your chemotherapy nurse or doctor if you have any of these symptoms. If you have this reaction, you will normally need an injection before each treatment.

- **Delayed or persistent diarrhoea**

If the diarrhoea starts more than 24 hours after your chemotherapy, you should immediately take the anti-diarrhoea treatment that the doctor prescribed for you. This is the loperamide dispensed by The Christie pharmacy. Follow your doctor's instructions exactly: take 2 loperamide capsules as soon as the first liquid stool occurs. Then take one capsule with each liquid loose stool. (The maximum dose of loperamide in a day is 8 capsules).

If the diarrhoea persists for more than 24 hours despite the loperamide you should start taking the ciprofloxacin antibiotic given to you on your first visit. Make sure you complete the 5-day course of the antibiotic. Contact The Christie if the diarrhoea persists. You can continue to take the loperamide.

You must tell your doctor if...

- you have a temperature/fever as well as diarrhoea
- you have nausea/vomiting as well as diarrhoea
- you still have diarrhoea 48 hours after starting the diarrhoea treatment

- **Rash**

Temozolomide may cause an itchy rash. If this happens, stop taking your tablets and let your hospital doctor know.

- **Sore mouth**

Your mouth may become sore or dry, or you may notice small mouth ulcers during this treatment. Drinking plenty of fluids and cleaning your teeth regularly and gently with a soft toothbrush can help to reduce the risk of this happening. We can prescribe a mouthwash for you to use during treatment. You can dilute this with water if your mouth is sore. For ask your doctor or nurse for further advice. There is also general mouth care information in the chemotherapy booklet. If you continue to have a sore mouth, please contact The Christie Hotline. Occasionally during treatment you may experience a strange taste sometimes described as metallic or bitter. A strongly flavoured sweet or mint will help to disguise this.

- **Skin changes**

Sometimes as a result of the chemotherapy, your skin may appear darker in colour or lightly tanned, especially around the joints. This is known as hyperpigmentation. The skin will return to normal when treatment is finished. Some chemotherapy can make your skin more sensitive to the sun than usual. Sit in the shade, avoid too much sun and use a sunblock cream. Asian and African-Caribbean people may develop noticeable light patches on their skin.

- **Liver or kidney problems**

Temozolomide can rarely affect liver and kidney function. Your kidney and liver function will be monitored every time you come to the clinic for your treatment.

Serious and potentially life threatening side effects

In a small proportion of patients chemotherapy can result in very severe side effects which may rarely result in death. The team caring for you will discuss the risk of these side effects with you.

- **Herbal medicine**

Some herbal medicine including St John's Wort can affect the chemotherapy. You should let your doctor or nurse know if you are taking any herbal medication, complementary or alternative medicines, including vitamins, minerals and medicines purchased over-the-counter.

Sex, contraception & fertility

Protecting your partner and contraception: We recommend that you or your partner use a condom during sexual intercourse while you are having the course of chemotherapy. Chemotherapy is dangerous to unborn babies and this will also protect you and your partner from any chemotherapy drugs that may be present in semen and in the vagina. If you suspect that you may be pregnant please tell your doctor immediately.

Fertility: This chemotherapy may affect your ability to have children. Your doctor or nurse should have discussed this with you. If not, please ask them before you start treatment.

Late side effects

Some side effects may become evident only after a number of years. In reaching any decision with you about treatment, the potential benefit you receive from treatment will be weighed against the risks of serious long term side effects to the heart, lungs, kidneys and bone marrow. With some drugs there is also a small but definite risk of developing another cancer. If any of these problems specifically applies to you, the doctor will discuss these with you and note this on your consent form.

Contacts

If you have any general questions or concerns about your treatment, please ring the area where you are having treatment:


Administrative enquiries	0161 918 7606/7610
Chemotherapy nurse:	0161 918 7171
Clinical trials unit	0161 918 7663
Sarcoma Clinical Nurse Specialist	0161 446 3094
Sarcoma Research Office	0161 918 7355 / 7098
Palatine Treatment Ward	0161 446 3960

For advice ring The Christie Hotline on 0161 446 3658 (24 hours)

Your consultant is: _____

Your hospital number is: _____

Your key worker is: _____



The Christie Patient Information Service July 2014
CHR/CT/711/02.02.10 version 3 Review date: July 2017

Details of the sources used are available, please contact Patient.Information@christie.nhs.uk