

Pemetrexed and carboplatin

The possible benefits of treatment vary; for some people this chemotherapy may reduce the risk of the cancer coming back, for others it may control the cancer and its symptoms. Your doctor will explain to you whether you will receive chemotherapy or another type of treatment, or a combination of both. They will be happy to answer any questions you have about your treatment. You may find the booklet 'Chemotherapy, a guide' helpful, which gives general information on chemotherapy and side effects.

Your treatment

Your doctor has prescribed for you a treatment which includes the chemotherapy pemetrexed (Alimta®) and carboplatin given into the vein via a drip.

The treatment consists of the following:

Day 1 Pemetrexed (Alimta®) and carboplatin by a drip over 1-2 hours.

Day 15 No treatment, however we may ask you to attend for a clinic visit.

Day 21 Restart with the next cycle (day 1).

The treatment is repeated every 3 weeks for a total of 4 to 6 cycles.

In some cases pemetrexed may be continued beyond this point as a maintenance treatment.

You will have a routine blood test before the start of every cycle of treatment.

Occasionally we may not be able to go ahead with your treatment until your blood counts are back to a safe level. If this happens, your chemotherapy may be delayed a week.

IMPORTANT: You will need to take some medications before starting the treatment.

Details of these are below. Remember to take your dexamethasone tablets twice daily for 3 days (take the first dose in the morning with breakfast and take the second dose at lunch-time). **Start taking the dexamethasone the day before your chemotherapy.** This is important – dexamethasone helps to prevent allergic reactions to the chemotherapy. If you forget, tell your chemotherapy nurse before you have treatment.

- **Folic acid tablets** need to be taken once daily, continuously throughout treatment. This will start at least 5 days before the chemotherapy. If you are running out of these tablets at any point during your treatment, please let your nurse or doctor know.
- You will also have **Vitamin B12** injections. These will be given on the first day of chemotherapy treatment for cycle 1 and cycle 4, then every 3 cycles thereafter.
- Anti-inflammatory drugs such as ibuprofen, diclofenac etc should be discontinued. We strongly advise current smokers **to stop smoking while having this treatment.** If you would like help with stopping smoking, please let your doctor or nurse know and they can arrange support for you.

This treatment can have serious or possibly life-threatening side effects. It is very important that you report side effects straight away. Don't delay, if you feel unwell, please ring The Christie Hotline on **0161 446 3658**. The lines are open 24 hours a day.



Kidney function

It is vital to measure how your kidneys are working before and during your treatment and this is done by checking your blood results, or we may ask you to have a GFR test (nuclear medicine test) which involves an extra visit to hospital for approximately 4 hours.

Dexamethasone

Remember to take your dexamethasone tablets twice daily for 3 days (or according to your doctors instruction). Start taking the dexamethasone the day before your chemotherapy. This is important - dexamethasone helps to prevent allergic reactions to chemotherapy. If you forget, tell your chemotherapy nurse before you have treatment.

Please note:

- Your **dexamethasone tablets** should be taken with food as they may cause indigestion. They may also increase your appetite. You should take them earlier in the day, for example with breakfast and lunch, since they can make you feel more alert and prevent sleep.
- Dexamethasone tablets can **alter your mood** causing you to feel low, irritable or agitated. Please tell the doctor at your next clinic if this is a problem.
- Some people may have an **immediate allergic reaction** to the chemotherapy. You may feel short of breath, develop a rash on your body and have red flushes. This is temporary but call your nurse immediately. The nurse can give you an injection to stop this.

Blood sugar

Steroids can cause your blood sugar to rise. If you are diabetic then you may need to increase the dose of any diabetic medication you take (insulin or tablets). You should discuss this with your doctor before starting the chemotherapy.

Increased risk of serious infection

You are vulnerable to infection while you are having chemotherapy. Minor infections can become life-threatening in a matter of hours if left untreated. Symptoms of infection include fever, shivering, sweats, sore throat, diarrhoea, discomfort when you pass urine, cough or breathlessness. We recommend that you use a digital thermometer so you can check your temperature. You can buy one from your local chemist. You may be given injections or antibiotics to help reduce the risk of having a serious infection.

If you feel unwell, you have symptoms of an infection or your temperature is 37.5°C or above, or below 36°C contact The Christie Hotline straight away.

Possible side effects

Chemotherapy can cause many different side effects. Some are more likely to occur than others. Everyone is different and not everyone gets all the side effects. Most side effects are usually temporary, but in some rare cases they can be life-threatening. It is important to tell your hospital doctor or nurse about any side effects so they can be monitored and, where possible, treated.

Common side effects (More than 1 in 10)

• Anaemia (low number of red blood cells)

Whilst having your treatment you may become anaemic. This may make you feel tired and breathless. Let your doctor or nurse know if you are affected by these symptoms. In some instances, you may require a blood transfusion.

- **Bruising or bleeding**

This treatment can reduce the production of platelets which help the blood clot. Tell your doctor if you have any unexplained bruising or bleeding, such as nosebleeds, blood-spots or rashes on the skin, and bleeding gums. You may need a platelet transfusion.

- **Nausea and vomiting (sickness)**

The severity of this varies from person to person. Anti-sickness medication may be given along with your chemotherapy to prevent this. You may also be given anti-sickness tablets to take at home. If you continue to feel or be sick, contact your GP or this hospital, because your anti-sickness medication may need to be changed or increased.

- **Lethargy**

Some chemotherapy may make you feel tired and lacking in energy. It can be frustrating when you feel unable to cope with routine tasks. If you do feel tired take rest and get help with household chores. If necessary take time off work. Gentle exercise such as a daily walk can be beneficial. Tiredness should not be profound. If you are unable to get up and get dressed by yourself contact the hotline for advice.

- **Loss of appetite**

If you experience a loss of appetite, please be sure to tell your doctor or nurse at your next hospital visit.

- **Sore mouth**

Your mouth may become sore or dry, or you may notice small mouth ulcers during this treatment. Drinking plenty of fluids and cleaning your teeth regularly and gently with a soft toothbrush can help to reduce the risk of this happening. We can prescribe a mouthwash for you to use during treatment. You can dilute this with water if your mouth is sore. Ask your doctor or nurse for further advice. There is also general mouth care information in the chemotherapy booklet. If you continue to have a sore mouth, please contact The Christie Hotline on **0161 446 3658**.

- **Diarrhoea**

If this becomes a problem while you are having treatment, anti-diarrhoea tablets can be bought from a pharmacy or prescribed by your GP for a temporary period until this is resolved. If the problem persists contact The Christie. If you develop severe diarrhoea it is important to contact The Christie Hotline on **0161 446 3658** straight away as this may be a sign of a serious infection. Don't delay!

- **Constipation**

Try to drink plenty of fluids and eat foods high in fibre. Report this to your hospital doctor who may prescribe a suitable laxative. Ask the staff for a copy of 'Eating - help yourself' which has useful ideas about diet when you are having treatment.

- **Skin rash**

You may develop a skin rash, this is usually mild and easily treated. Please tell your doctor on your next visit.

- **Kidney damage**

There is a small risk of damage to the kidneys from the carboplatin and pemetrexed. Your medical team will continue to monitor your bloods at each visit and will discuss with you if there are any concerns. It is important to drink plenty of water/fluids whilst on treatment.

- **Possible hair thinning**

You may lose your hair although this is not usually total hair loss. The hair falls out gradually 10 to 14 days following your treatment. The timescale varies from person to person. Please remember that this is temporary side effect and your hair will grow back when your treatment is completed. Very rarely, hair loss can be permanent.

The cancer information centre offers a coping with hair loss service to all patients where support, information and advice will be given. Drop in, contact **0161 446 8100** or email

the-christie.informationcentre@nhs.net

Information about the wig service can also be found here and vouchers for wigs can also be obtained for eligible patients. The wig room also provides a drop in service, please see The Christie leaflet 'The wig fitting service' for further information.

Less common side effects (Less than 1 in 10)

• Immediate allergic reactions

Allergic reactions to carboplatin are uncommon, but please ask the staff for help immediately if you notice any of the following: fevers and chills; back pain; shortness of breath; headaches and swelling of the face may occur during the time the drug is being given. If this happens please tell the staff straight away. Your doctor may prescribe medication that can help to reduce these side-effects and, if necessary, we can give you this before your next treatment.

• Tinnitus and high-frequency hearing loss

You may develop tinnitus (ringing in the ears). This sensation should subside when your treatment finishes. Please tell your doctor if this occurs. High frequency hearing loss / difficulty hearing high pitched sounds, can also occur with this chemotherapy. This may be permanent.

• Tingling and numbness in the fingers or toes

Usually only mild and temporary. Please report these symptoms to your doctor on your next hospital visit. On rare occasions, this may be permanent.

• Strange taste

Occasionally during treatment you may experience a strange taste, sometimes described as metallic or bitter. A strongly flavoured sweet or mint will help to disguise this.

Rare side effects (less than 1 in 100)

• Chest pain (Warning!)

If you experience chest pain, please tell your doctor or nurse so you can discuss treatment options. In an emergency you should go straight to your nearest accident and emergency department.

Serious and potentially life threatening side effects

In a small proportion of patients chemotherapy can result in very severe side effects which may rarely result in death. The team caring for you will discuss the risk of these side effects with you.

Other medicines

Please ask your doctor at The Christie for advice about any other medication you are taking, including non-prescribed medicines, complementary therapies and herbal medicines.

Sex, contraception and fertility

Protecting your partner and contraception: We recommend that you or your partner use a condom during sexual intercourse while you are having the course of chemotherapy. Chemotherapy is dangerous to unborn babies and this will also protect you and your partner from any chemotherapy drugs that may be present in semen and in the vagina. If you suspect that you may be pregnant please tell your doctor immediately.

Fertility: This chemotherapy may affect your ability to have children. Your doctor or nurse should have discussed this with you. If not, please ask them before you start treatment.

For female patients only

Menopausal symptoms

When the ovaries stop working due to chemotherapy or during a natural menopause most women experience symptoms such as hot flushes, sweats (night and day) and vaginal dryness. These hormonal changes can make the vagina feel as though it has shrunk and become less easy to stretch. This is called vaginal atrophy and can result in discomfort, pain on sexual intercourse, itching and recurrent urine infections. If your ovaries don't start to work again the vaginal symptoms can be permanent, although the flushes and sweats tend to reduce and stop over a small number of years. Some women who have already gone through menopause may notice their symptoms worsening for a time after chemotherapy.

The vaginal symptoms can start early and the longer they are left the harder they can be to treat. Please contact your specialist nurse either in clinic or by phone when the symptoms first develop if you would like help. Symptoms can be managed in several ways including gels, essential oil pessaries and sometimes local oestrogen replacement. You may also find it helpful to visit the page 'Managing menopausal symptoms after cancer treatment' on the Macmillan Cancer Support website:

www.macmillan.org.uk

Late side effects

Some side effects may become evident only after a number of years. In reaching any decision with you about treatment, the potential benefit you receive from treatment will be weighed against the risks of serious long term side effects to the heart, lungs, kidneys and bone marrow. With some drugs there is also a small but definite risk of developing another cancer. If any of these problems specifically applies to you, the doctor will discuss these with you and note this on your consent form.

Contacts

If you have any general questions or concerns about your treatment, please ring the area where you are having treatment:

Administration enquiries - **0161 918 7606/7610**

Chemotherapy nurse - **0161 918 7171**

Clinical trials unit - **0161 918 7663**

Lung cancer nurse team - **0161 918 2595**

Your doctor's secretary **0161**

For urgent advice ring The Christie Hotline on **0161 446 3658** (24 hours a day, 7 days a week)

Your consultant is:

Your hospital number is:

Your key worker is:

© 2024 The Christie NHS Foundation Trust. This document may be copied for use within the NHS only on the condition that The Christie NHS Foundation Trust is acknowledged as the creator.

If you need information in a different format, such as easy read, large print, BSL, braille, email, SMS text or other communication support, please tell your ward or clinic nurse.

The Christie is committed to producing high quality, evidence based information for patients. Our patient information adheres to the principles and quality statements of the Information Standard. If you would like to have details about the sources used please contact **the-christie.patient.information@nhs.net**

For information and advice visit the cancer information centres at Withington, Oldham, Salford or Macclesfield. Opening times can vary, please check before making a special journey.



Contact The Christie Hotline for
urgent support and specialist advice
The Christie Hotline: 0161 446 3658
Open 24 hours a day, 7 days a week