

## Patient guide to R-CODOX M

Day 1	Rituximab	Given as an infusion. The first dose is started slowly and can take a few hours. If no reactions subsequent infusion given over 60 mins.
	Doxorubicin	Given as injection (bolus) diluted through a saline drip over 5-10 mins.
	Cyclophosphamide	Given as bolus or infusion over 5-10 mins.
	Vincristine	Given as an infusion over 5-10 mins.
Day 2	Cyclophosphamide	Given as bolus or infusion over 5-10 mins.
(or nearest Tuesday/Friday)	Intrathecal Cytarabine	Injection into spinal fluid. Performed by specially trained doctor. Administered on the haematology day unit. All intrathecal injections will require platelet count of 50 or higher and normal clotting levels, these will be checked on the day.
Days 3 and 4	Cyclophosphamide	Given as bolus or infusion over 5-10 mins.
Day 5	Cyclophosphamide	Given as bolus or infusion over 5-10 mins.
(or nearest Tuesday/Friday)	Intrathecal Cytarabine	See day 2 for Intrathecal chemotherapy notes.
Day 6 & 7	No chemotherapy	You may go home for these days if you are well and the blood count satisfactory.
Day 8	Rituximab	Given as an infusion over 60 mins (if no previous infusion reaction).
	Vincristine	Given as bolus over 5-10 mins.
Day 9	No chemotherapy	You may go home for this day if you are well and the blood count satisfactory.
Day 10	I.V. Methotrexate	Given as an infusion for 1 hour, then followed by a 23 hour infusion. You will also have fluids running alongside the chemotherapy. Folinic acid rescue will be given, initially by injection then by mouth. Methotrexate blood levels will be checked at specific times.
Days 11 to 14	No chemotherapy	
Day 15	Intrathecal	See day 2 Intrathecal chemotherapy notes.
(or nearest Tuesday/Friday)	Methotrexate	
Day 16 until blood count recovery	No chemotherapy	Waiting for blood counts to recover before starting next cycle of chemotherapy.





Contact The Christie Hotline for urgent support and specialist advice

The Christie Hotline: 0161 446 3658

Open 24 hours a day, 7 days a week