

Carboplatin

The possible benefits of treatment vary; for some people this chemotherapy may reduce the risk of the cancer coming back, for others it may control the cancer and its symptoms. Your doctor will explain to you whether you will receive chemotherapy or another type of treatment, or a combination of both. Your doctor or nurse will be happy to answer any questions you have about your treatment. You will find it useful to refer to the booklet 'Chemotherapy, a guide' which gives general information on chemotherapy and side effects.

Your treatment

Your doctor or nurse clinician has prescribed for you treatment with carboplatin chemotherapy. The chemotherapy is given into a vein via a drip over 1 hour. The treatment is repeated every 3 - 4 weeks for total of 6 cycles.

You will have a routine blood test before the start of each cycle of treatment. Occasionally, we may not be able to go ahead with your treatment until your blood counts are back to a safe level. If this happens, your chemotherapy may be delayed a week.

Kidney function. It is important to monitor how your kidneys are working while you are having treatment. We can do this by checking your blood results or we may ask you to attend for a 'GFR' test.

Carboplatin chemotherapy may be given in combination with another chemotherapy.

This treatment can have serious or possibly life-threatening side effects. It is very important that you report side effects straight away. Don't delay, if you feel unwell, please ring The Christie Hotline on **0161 446 3658**. The lines are open 24 hours a day.

Increased risk of serious infection (Warning!)

You are vulnerable to infection while you are having chemotherapy. Minor infections can become life-threatening in a matter of hours if left untreated. Symptoms of infection include fever, shivering, sweats, sore throat, diarrhoea, discomfort when you pass urine, cough or breathlessness. We recommend that you use a digital thermometer so you can check your temperature. You can buy one from your local chemist.

If you feel unwell, you have symptoms of an infection or your temperature is 37.5°C or above, or below 36°C contact The Christie Hotline straight away.



243 Carboplatin

Common side effects (more than 1 in 10)

Anaemia (low number of red blood cells) (Warning!)

While having this treatment you may become anaemic. This may make you feel tired and breathless. Let your doctor or nurse know if these symptoms are a problem. You may need a blood transfusion.

Nausea and vomiting (sickness)

The severity of this varies from person to person. Anti-sickness medication may be given along with your chemotherapy to prevent this. You will also be given anti-sickness tablets to take at home. If you continue to feel or be sick, contact your GP or this hospital, because your anti-sickness medication may need to be changed or increased.

Lethargy

Some chemotherapy may make you feel tired and lacking in energy. It can be frustrating when you feel unable to cope with routine tasks. If you do feel tired, take rest and get help with household chores. If necessary, take time off work. Gentle exercise such as walking can be beneficial.

• Diarrhoea

If this becomes a problem while you are having treatment, anti-diarrhoea tablets can be bought from a pharmacy or prescribed by your GP for a temporary period until this is resolved. If the problem persists contact The Christie. Ask the staff for a copy of 'Eating: help yourself 'which has some useful ideas about diet when you are having treatment.

Constipation

Try to drink plenty of fluids. Report this to your hospital doctor or nurse who can advise you regarding diet and who may prescribe a suitable laxative. Ask the staff for a copy of Eating: help yourself which has useful ideas about diet when you are having treatment.

Hair thinning

Some hair loss may occur during treatment, including body and facial hair. It is advisable to avoid perms, colours, use of hot brushes and vigorous frequent washing that could increase hair loss. Please remember that this is a temporary side effect and your hair will grow back when your treatment is completed. Very rarely, hair loss may be permanent. If you would like an appointment with the wig service, this can be arranged for you by visiting the cancer information centre. It is a good idea to get your wig before you lose a lot of hair which you can then match to your natural colour. Ask the staff for a copy of the 'Wig fitting service at The Christie'.

The Maggie's Centre runs a Talking Heads hair loss support workshop for anyone who is anticipating or experiencing hair loss (both men and women). These sessions cover the practicalities of hair loss as well as offering support with its emotional impact. Contact Maggie's on **0161 641 4848** or email manchester@maggiescentres.org

Uncommon side effects (less than 1 in 10)

• Bruising or bleeding (Warning!)

This treatment can reduce the production of platelets which help the blood clot. Let your doctor know if you have any unexplained bruising or bleeding, such as nosebleeds, bloodspots or rashes on the skin, and bleeding gums. You may need a platelet transfusion.

Kidney function

It is important to monitor how your kidneys are working while you are having treatment. We do this through routine blood tests, or GFR test (most accurate test of kidney function). It is important to drink plenty of fluids (at least 8 cups) the day before and for a few days after chemotherapy.

243 Carboplatin Page 2 of 4

Rare side effects (less than 1 in 100)

Tingling and numbness in the fingers or toes

This is uncommon. It is usually only mild and temporary but it can gradually build up during the treatment course and occasionally become permanent. Please report these symptoms to your doctor on your next hospital visit.

Tinnitus and high frequency hearing loss

You may develop tinnitus (ringing in the ears), this sensation should subside when your treatment finishes. High frequency hearing loss can occur with this chemotherapy. Rarely, this may be permanent.

Immediate allergic reactions (Warning!)

Allergic reactions to carboplatin are uncommon. Please ask the staff for help *immediately* if you notice any of the following: fevers and chills, back pain, shortness of breath, headaches and swelling of the face may occur during the time the drug is being given. If this happens please tell the staff straight away. Your doctor may prescribe further medication that can help to reduce these side effects.

Extravasation

Extravasation is when chemotherapy leaks outside the vein. If you develop redness, soreness or pain at the injection site **at any time** please let us know straight away. If you feel pain, tell your doctor or nurse as they can slow the drip to reduce the reaction.

Serious and potentially life threatening side effects

In a small proportion of patients chemotherapy can result in very severe side effects which may rarely result in death. The team caring for you will discuss the risk of these side effects with you.

Other medicines

Some medicines can be harmful to take when you are having chemotherapy. Let your doctor know about any medications you are taking, including non-prescribed medicines such as complementary therapies and herbal remedies.

Sex, contraception and fertility

Protecting your partner and contraception: We recommend that you or your partner use a condom during sexual intercourse while you are having the course of chemotherapy. Chemotherapy is dangerous to unborn babies and this will also protect you and your partner from any chemotherapy drugs that may be present in semen and in the vagina. If you suspect that you may be pregnant please tell your doctor immediately.

Fertility: This chemotherapy may affect your ability to have children. Your doctor or nurse should have discussed this with you. If not, please ask them before you start treatment.

Loss of periods: Due to the effects of chemotherapy on the ovaries you may find that your periods become irregular or may eventually stop. In younger women this may be temporary, but if you are closer to your menopause it may be permanent. This will result in hot flushes, sweats and vaginal dryness.

Late side effects


Some side effects may become evident only after a number of years. In reaching any decision with you about treatment, the potential benefit you receive from treatment will be weighed against the risks of serious long term side effects to the heart, lungs, kidneys and bone marrow. With some drugs there is also a small but definite risk of developing another cancer. If any of these problems specifically applies to you, the doctor will discuss these with you and note this on your consent form.

Contacts

If you have any general questions or concerns about your treatment, please ring the area where you are having treatment:

Administration enquiries - 0161 918 7606/7610 Chemotherapy nurse - 0161 918 7171 Clinical trials unit - 0161 918 7663

For advice ring the Christie Hotline on U161 446 3658 (24 hours
Your consultant is:
Your hospital number is:
Your key worker is:



If you need information in a different format, such as easy read, large print, BSL, braille, email, SMS text or other communication support, please tell your ward or clinic nurse.

The Christie is committed to producing high quality, evidence based information for patients. Our patient information adheres to the principles and quality statements of the Information Standard. If you would like to have details about the sources used please contact **patient.information@christie.nhs.uk**

For information and advice visit the cancer information centres at Withington, Oldham or Salford. Opening times can vary, please check before making a special journey.



Contact The Christie Hotline for urgent support and specialist advice

The Christie Hotline: 0161 446 3658 Open 24 hours a day, 7 days a week

July 2020 – Review July 2023 CHR/CT/243/12.01.04 Version 10 The Christie Patient Information Service Tel: 0161 446 3000 www.christie.nhs.uk